

Prof. David Draper  
Department of Statistics  
University of California, Santa Cruz

## AMS 131: Quiz 9

Name: \_\_\_\_\_

You're about to take an IID sample  $(X_1, \dots, X_n)$  from a distribution with variance  $V(X_i) = \sigma^2$  that exists and is finite, which implies that the mean  $E(X_i) = \mu$  also exists and is finite. The purpose of the sampling is to use the sample mean  $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$  as an estimator of  $\mu$ , and you're wondering what value you should use for the sample size  $n$ . We've seen in class that *Chebyshev's Inequality* can help when no other details about the distribution of the  $X_i$  are available: if  $Y$  is any random variable whose variance  $V(Y)$  exists, this inequality states that for any  $t \geq 0$

$$P(|Y - E(Y)| \geq t) \leq \frac{V(Y)}{t^2}. \quad (1)$$

- (a) Using basic facts about  $E(\bar{X}_n)$  and  $V(\bar{X}_n)$ , show that inequality (1) implies, in the random sampling problem considered here, that for any  $k > 0$

$$P(|\bar{X}_n - \mu| < k\sigma) \geq 1 - \frac{1}{nk^2}, \quad (2)$$

and show further that, if we want the probability in (2) to be at least  $(1 - \alpha)$  for some  $0 < \alpha < 1$ , we should choose

$$n_{Chebyshev} \geq \frac{1}{\alpha k^2}. \quad (3)$$

I also mentioned in class that Chebyshev's Inequality can be quite conservative. Let's quantify this: suppose for the rest of the problem that  $(X_i | \mu, \sigma^2) \stackrel{IID}{\sim} N(\mu, \sigma^2)$ . Then we've seen in class that  $\bar{X}_n$  also follows a Normal distribution, with mean  $\mu$  and standard error  $SE(\bar{X}_n) = \frac{\sigma}{\sqrt{n}}$ . As usual let  $\Phi(x)$  be the standard Normal CDF; in other words,  $\Phi(x)$  is the area to the left of  $x$  under the standard Normal PDF.

- (b) Sketch the PDF of  $\bar{X}_n$ , shading in the area corresponding to  $(*) = (|\bar{X}_n - \mu| < k\sigma)$  and identifying the places on both the raw-units and standard-units axes corresponding to the endpoints of  $(*)$ .

- (c) Using a basic fact about  $\Phi(x)$  and your sketch in part (b), show that under the Normality assumption for the  $X_i$ ,

$$P(|\bar{X}_n - \mu| < k\sigma) \geq 2\Phi(k\sqrt{n}) - 1. \quad (4)$$

- (d) Set the probability on the right-hand side of the inequality in (4) equal to  $(1 - \alpha)$  and solve for  $n$ , thereby showing that under the Normality assumption the required sample size corresponding to  $n_{Chebyshev}$  in equation (3) above is

$$n_{Normality} \geq \frac{[\Phi^{-1}(1 - \frac{\alpha}{2})]^2}{k^2}, \quad (5)$$

in which (as usual)  $\Phi^{-1}(p)$  (for  $0 < p < 1$ ) is the inverse CDF (quantile function) for the standard Normal distribution.

- (e) Using the table on page 861 of Degroot and Schervish or (better) an online inverse Normal CDF calculator (e.g., there's one provided by [Wolfram Alpha](#)), complete the rest of the table below.

$\alpha$	$\frac{1}{\alpha}$	$[\Phi^{-1}(1 - \frac{\alpha}{2})]^2$
0.1	10	2.7
0.05	20	
0.01		6.6
0.005		
0.001	1000	

- (f) If the data values  $X_i$  really did come from a Normal distribution, would you describe the Chebyshev Inequality sample size calculation as highly conservative, not too conservative, or in between? Explain briefly.